



Patient and Doctor Discussion Guide

Ask yourself—Is RUCONEST right for me?

If you answer **YES** to any of the questions below, you may be ready for RUCONEST.

RUCONEST is a plasma-free C1 esterase inhibitor (C1-INH) therapy for the treatment of acute attacks in adults and adolescents with hereditary angioedema (HAE). Effectiveness in clinical studies was not established in HAE patients with laryngeal attacks.

The serious adverse reaction reported in clinical studies of RUCONEST was anaphylaxis. The most common adverse reactions reported were headache, nausea, and diarrhea.

Are you interested in a therapy that can treat an attack in just one dose?

Yes No

Is it important to you to be as prepared as possible for an HAE attack?

Yes No

Are you interested in a plasma-free therapy for HAE that treats the problem at its source?

Yes No

Ask your doctor about RUCONEST, a plasma-free C1-INH therapy.

To learn more about RUCONEST, call **1-855-613-4HAE (4423)** or visit www.ruconest.com today.

Please see accompanying Important Safety Information on next page.

 **RUCONEST**[®]
C1 esterase inhibitor (recombinant)

IMPORTANT SAFETY INFORMATION

RUCONEST® (C1 esterase inhibitor [recombinant]) is not for everyone. Do not take RUCONEST if you have a known history of allergy to rabbits or products from rabbits. Do not take RUCONEST if you have a history of life-threatening immediate allergic reactions to C1 esterase inhibitor preparations, including anaphylaxis.

If you experience hives, pale red, raised, itchy bumps (urticaria), tightness of the chest, wheezing, low blood pressure (hypotension), and/or anaphylaxis during or after injection of RUCONEST, discontinue RUCONEST and immediately contact your doctor. These may be signs and symptoms of allergic reactions.

Products similar to RUCONEST have been associated with thromboembolic events. Before taking RUCONEST, please notify your doctor if you have an indwelling venous catheter/access device, history of blood clot (thrombosis), been told you have thickening of the walls of your arteries (atherosclerosis), use oral contraceptives (i.e. estrogen or progesterone), are extremely overweight and have significant difficulty moving around.

If you are pregnant, planning to become pregnant, or nursing, talk to your healthcare provider before taking RUCONEST.

The most common RUCONEST side effects in clinical studies include: headache, nausea, and diarrhea. Serious side effect anaphylaxis has been reported in RUCONEST clinical studies.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

For product information, adverse event reports, and product complaint reports, please contact Pharming U.S. toll-free:

(844) 474-2764 Main corporate number

(800) 930-5221 Pharmacovigilance, product complaints, medical information

Please see accompanying full Prescribing Information, including Patient Product Information [here](#), or visit www.ruconest.com.