

# **Patient Enrollment Form**

Fax completed forms to: 1-855-423-5757



## **Getting Started**

## Step 1:

### Fill out all 3 pages of the Enrollment Form

#### Page 1

### Patient to read and sign the Consent Form

**NOTE**: Patient signature on Consent Form is required to access RUCONEST Solutions support, including RPA services, copay assistance, and any free goods program, including StarterRx. If consent is not submitted with the enrollment form, RUCONEST Solutions will work with the patient to obtain consent.

### Pages 2 and 3

<u>Provider to fill out and sign the Enrollment Form</u> including a copy of the patient's insurance card

**NOTE**: The Enrollment Form provides prescription for both Commercial and Free Goods Programs

## Step 2:

Submit Pages 1 through 3 of the Enrollment Form to RUCONEST Solutions, along with the following documentation:

- Copy of recent labs related to HAE
- Patient's current weight and full medication list, including previous and current HAE therapies and any known drug or other allergies
- Clinical notes documenting patient signs, symptoms, and manifestations of HAE
- Any <u>swell log or diary details</u> of recent frequency, severity, and duration of acute HAE attacks
- Any additional clinical information pertaining to patient's clinical history
- · Documentation of other therapies used to treat symptoms of HAE

This requested documentation will help RUCONEST Solutions to support your office with coverage authorizations when allowed by an insurance company. There may be occasions where the insurer will request additional documentation and/or mandate that your office submit the coverage requests. If this is the case, your office will be informed on a subsequent fax or phone call from the RUCONEST Solutions support team.

## Step 3:

Let your patient know you are sending in a referral for them and that RUCONEST Solutions will be calling them





Questions? Call 855-613-4423 between 8 am-8 pm ET M-F for additional assistance.

**RUCONEST Solutions: 1-855-613-4423** 

### **INDICATIONS AND USAGE**

RUCONEST® (C1 esterase inhibitor [recombinant]) is indicated for the treatment of acute attacks in adult and adolescent patients with hereditary angioedema (HAE). Effectiveness in clinical studies was not established in HAE patients with laryngeal attacks.

#### IMPORTANT SAFETY INFORMATION

RUCONEST is contraindicated in patients with a history of allergy to rabbits or rabbit-derived products and for patients with a history of life-threatening immediate hypersensitivity reactions, including anaphylaxis, to C1 esterase inhibitor (C1-INH) preparations.

Monitor patients for early signs of allergic or hypersensitivity reactions (including hives, generalized urticaria, tightness of the chest, wheezing, hypotension, and/or anaphylaxis). If symptoms occur, discontinue RUCONEST and administer appropriate treatment.

Serious arterial and venous thromboembolic (TE) events have been reported with plasma-derived C1-INH products. Risk factors may include the presence of an indwelling venous catheter/access device, prior history of thrombosis, underlying atherosclerosis, use of oral contraceptives or certain androgens, morbid obesity, and immobility. Monitor patients with known risk factors for TE events during and after RUCONEST administration.

Appropriately trained patients may self-administer RUCONEST upon recognition of an HAE attack. Advise patients to seek medical attention if progress of any attack makes them unable to properly prepare or administer a dose of RUCONEST. No more than 2 doses should be administered within a 24-hour period.

The serious adverse reaction reported in clinical trials was anaphylaxis. The most common adverse reactions (incidence ≥2%) were headache, nausea, and diarrhea.

Before prescribing RUCONEST, please read the accompanying full Prescribing Information or go to www.ruconest.com



# **Patient Enrollment Form**

**Fax completed forms to: 1-855-423-5757** 



## **Patient Consent Form**

Patient Name:	DOB:
Patient Email:	Patient Phone (Cell):
Emergency Contact Name:	Relationship to Patient:
Emergency Contact Phone:	_
Consent to Share Health Information: By signing this Consent, I aurhome care nurse educators, my health plan(s) providing medical care the RUCONEST to disclose to RUCONEST Solutions ("Program") open Pharming Healthcare, health information relating to my medical cond my healthcare provider, including physicians and home care nurse ed prescription coverage, and my pharmacy(ies) providing the RUCONEST condition, treatment, and insurance coverage from the Program. I authorized (and related information and materials) related to any of Pharmsurance coverage, prescription fulfillment, online support, financial services; and (ii) information about Pharming Healthcare's products, use my health information to conduct data analytics, market research information has been disclosed to Pharming Healthcare, I understand However, Pharming Healthcare agrees to protect my health information this Consent or as required by law or regulations. I understand that Pharming Healthcare in exchange for the health information and/or for that I may refuse to sign this Consent. I further understand that my trinsurance enrollment, or eligibility for insurance benefits are not concount sign it or later cancel it, I will not be able to receive Pharming Health at any time by calling (855) 613-4423. Canceling this Consent will end to Pharming Healthcare by my Healthcare Entities after they are notif by them pursuant to this Consent. Canceling this Consent will not affeinsurance. This Consent expires five (5) years from the date signed unless the signed unless that the pursuant to this Consent.	and prescription coverage, and my pharmacy(ies) providing rated by Pharming Healthcare and companies working with lition, treatment, and insurance coverage. I also authorize ucators, my health plan(s) providing medical care and ST to receive health information related to my medical chorize Pharming Healthcare to provide me with (i) support arming Healthcare's products, including but not limited to, assistance services, adherence, and other therapy support services, and programs. I understand that Pharming may n, and other internal business activities. Once my health that federal privacy laws no longer protect the information. On by using and disclosing it only for purposes authorized my pharmacy provider may receive remuneration from for any therapy support services provided to me. I understand reatment (including with a Pharming Healthcare product), ditioned upon my agreement to sign this Consent; but if I do lithcare's patient program support. I may cancel this consent I my consent to further disclosure of my health information fied of my cancellation but will not affect previous disclosures ect my ability to receive treatment, or my eligibility for health
Patient Support Services: I authorize the Program and its affiliated to any of Pharming Healthcare's products, including but not limited to assistance, financial assistance services, adherence, and other therap as well as any information or materials related to such services. I under Program are not employed by my healthcare professional. RUCONES email, fax, telephone call, text message (including calls and text mess prerecorded voice),* and other mutually agreed upon means. I also acconnection with the services and programs, including, without limitates.	o insurance coverage, prescription fulfillment, product by support services, relevant disease-related information, erstand that any personnel providing support as part of the T Solutions or Pharming Healthcare may contact me by mail, sages made with an automatic telephone dialing system or a uthorize Pharming Healthcare to use my health information in
<b>Opt-in for Other Resources:</b> By signing below, I authorize Pharming Healthcare, to contact me by mail, email, fax, text messaging,* and/o customer surveys, or occasionally for market research purposes. I uncondition of receiving any Pharming Healthcare medicine or Patient S or trade my personal data to any unrelated third party.	r telephone regarding other potential topics of interest to me, derstand that I am not required to provide this consent as a
☐ I would like to <b>opt out</b> of receiving other resources	
<b>Emergency Contact:</b> I authorize RUCONEST Solutions, my doctor, m me with access to RUCONEST to contact the emergency contact liste	
By signing below, I confirm that I have read and understand the Support Services above and agree to the terms.	e Consent to Share Health Information and Patient
Printed Patient/Legal Representative Name:	
Patient/Legal Representative Signature:	
If Legal Representative, Relationship to Patient:	

\*Data rates may apply.

Before prescribing RUCONEST, please read the accompanying full Prescribing Information or go to www.ruconest.com

RUCONEST Solutions: 1-855-613-4423



# **Patient Enrollment Form**

Fax completed forms to: 1-855-423-5757



Select which specialty pharmacy the patient currently uses (if known):

1. Prescription Accredo Health Grou	up CVS Caremark Orsini
Patient Name	ANCILLARY ORDERS: Dispense infusion supplies with each prescription.  Dispense: One (1) vial of Sterile Water for Injection 14 mL per 2100 IU vial of RUCONEST
Prescription: RUCONEST 2100 international units (IU)/vial injection (50 IU/kg), Max 4200 IU  DIRECTIONS: AdministerIU as a slow IV injection over 5 min PRN for attacks. No more than 2 doses within a 24-hour period	Flushing Orders  Normal saline 3 mL or 5 mL intravenous (peripheral line) or 10 mL intravenous (central line) before and after infusion, or as needed for line patency Heparin 10 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Heparin 100 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Flushing orders not needed
4 doses (8 vials) 8 doses (16 vials) Per Month 16 doses (32 vials) —doses (vials) Per Shipment	Anaphylaxis Order Specialty pharmacy to provide anaphylactic kit per provider protocol. Substitution permitted unless DAW specified
Refill 1 x year, unless noted otherwise 3 Refills 6 Refills 12 Refills Refills	Epinephrine #2 pack 0.15mg 0.3mg Refills: Inject IM as needed for anaphylaxis reaction. May repeat x 1 in 5 to 15 minutes if symptoms persist. SP to provide at first dispense.
Concurrent Medications	
Drug/Non-Drug Allergies	No Known Allergies
Substitution permitted Dispense as written	
PRESCRIBER	Print Date
I attest that I have a HIPAA form on file and RUCONEST Solutions is authorize	red to perform a benefits verification. I appoint Pharming Healthcare, Inc., RUCONEST is prescription described herein to the dispensing pharmacy by any means allowed under
2. Optional Prescription for StarterRx, Bridge-t	to-Therapy, and/or PAP Program
Patient Name	ANCHI ADV ODDEDC. Discourse infusion complice with each prescription
DOBPatient Weightkg/lbs Diagnosis: ICD-10-CM D84.1 (Defects in the complement system [HAE])	ANCILLARY ORDERS: Dispense infusion supplies with each prescription.  Dispense: One (1) vial of Sterile Water for Injection 14 mL per 2100 IU vial of RUCONEST
Diagnosis: ICD-10-CM D84.1 (Defects in the complement system [HAE]) Other:  Prescription: RUCONEST 2100 IU/vial injection (50 IU/kg), Max 4200 IU  DIRECTIONS: AdministerIU as a slow IV injection over 5 min PRN for attacks. No more than 2 doses	Dispense: One (1) vial of Sterile Water for Injection 14 mL per 2100 IU vial of
Diagnosis: ICD-10-CM D84.1 (Defects in the complement system [HAE]) Other:  Prescription: RUCONEST 2100 IU/vial injection (50 IU/kg), Max 4200 IU  DIRECTIONS: AdministerIU as a slow IV	Dispense: One (1) vial of Sterile Water for Injection 14 mL per 2100 iU vial of RUCONEST  Flushing Orders  Normal saline 3 mL or 5 mL intravenous (peripheral line) or 10 mL intravenous (central line) before and after infusion, or as needed for line patency Heparin 10 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Heparin 100 units/mL (#3mL or #5mL) use as a final flush for central line (QS)
Diagnosis: ICD-10-CM D84.1 (Defects in the complement system [HAE]) Other:  Prescription: RUCONEST 2100 IU/vial injection (50 IU/kg), Max 4200 IU  DIRECTIONS: AdministerIU as a slow IV injection over 5 min PRN for attacks. No more than 2 doses within a 24-hour period 2 doses (4 vials) Per Month	Dispense: One (1) vial of Sterile Water for Injection 14 mL per 2100 IU vial of RUCONEST  Flushing Orders  Normal saline 3 mL or 5 mL intravenous (peripheral line) or 10 mL intravenous (central line) before and after infusion, or as needed for line patency Heparin 10 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Heparin 100 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Flushing orders not needed  Concurrent Medications
Diagnosis: ICD-10-CM D84.1 (Defects in the complement system [HAE]) Other:  Prescription: RUCONEST 2100 IU/vial injection (50 IU/kg), Max 4200 IU  DIRECTIONS: AdministerIU as a slow IV injection over 5 min PRN for attacks. No more than 2 doses within a 24-hour period  2 doses (4 vials)	Dispense: One (1) vial of Sterile Water for Injection 14 mL per 2100 iU vial of RUCONEST  Flushing Orders  Normal saline 3 mL or 5 mL intravenous (peripheral line) or 10 mL intravenous (central line) before and after infusion, or as needed for line patency Heparin 10 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Heparin 100 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Flushing orders not needed  Concurrent Medications
Diagnosis: ICD-10-CM D84.1 (Defects in the complement system [HAE]) Other:  Prescription: RUCONEST 2100 IU/vial injection (50 IU/kg), Max 4200 IU  DIRECTIONS: AdministerIU as a slow IV injection over 5 min PRN for attacks. No more than 2 doses within a 24-hour period 2 doses (4 vials) Per Month	Dispense: One (1) vial of Sterile Water for Injection 14 mL per 2100 IU vial of RUCONEST  Flushing Orders  Normal saline 3 mL or 5 mL intravenous (peripheral line) or 10 mL intravenous (central line) before and after infusion, or as needed for line patency Heparin 10 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Heparin 100 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Flushing orders not needed  Concurrent Medications
Diagnosis: ICD-10-CM D84.1 (Defects in the complement system [HAE]) Other:  Prescription: RUCONEST 2100 IU/vial injection (50 IU/kg), Max 4200 IU  DIRECTIONS: AdministerIU as a slow IV injection over 5 min PRN for attacks. No more than 2 doses within a 24-hour period  2 doses (4 vials)	Dispense: One (1) vial of Sterile Water for Injection 14 mL per 2100 iU vial of RUCONEST  Flushing Orders  Normal saline 3 mL or 5 mL intravenous (peripheral line) or 10 mL intravenous (central line) before and after infusion, or as needed for line patency Heparin 10 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Heparin 100 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Flushing orders not needed  Concurrent Medications
Diagnosis: ICD-10-CM D84.1 (Defects in the complement system [HAE]) Other:  Prescription: RUCONEST 2100 IU/vial injection (50 IU/kg), Max 4200 IU  DIRECTIONS: Administer IU as a slow IV injection over 5 min PRN for attacks. No more than 2 doses within a 24-hour period  2 doses (4 vials)	Dispense: One (1) vial of Sterile Water for Injection 14 mL per 2100 iU vial of RUCONEST  Flushing Orders  Normal saline 3 mL or 5 mL intravenous (peripheral line) or 10 mL intravenous (central line) before and after infusion, or as needed for line patency Heparin 10 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Heparin 100 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Flushing orders not needed  Concurrent Medications
Diagnosis: ICD-10-CM D84.1 (Defects in the complement system [HAE]) Other:  Prescription: RUCONEST 2100 IU/vial injection (50 IU/kg), Max 4200 IU  DIRECTIONS: Administer IU as a slow IV injection over 5 min PRN for attacks. No more than 2 doses within a 24-hour period  2 doses (4 vials)	Dispense: One (1) vial of Sterile Water for Injection 14 mL per 2100 iU vial of RUCONEST  Flushing Orders  Normal saline 3 mL or 5 mL intravenous (peripheral line) or 10 mL intravenous (central line) before and after infusion, or as needed for line patency Heparin 10 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Heparin 100 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Flushing orders not needed  Concurrent Medications
Diagnosis: ICD-10-CM D84.1 (Defects in the complement system [HAE]) Other:  Prescription: RUCONEST 2100 IU/vial injection (50 IU/kg), Max 4200 IU  DIRECTIONS: Administer IU as a slow IV injection over 5 min PRN for attacks. No more than 2 doses within a 24-hour period  2 doses (4 vials)	Dispense: One (1) vial of Sterile Water for Injection 14 mL per 2100 IU vial of RUCONEST  Flushing Orders  Normal saline 3 mL or 5 mL intravenous (peripheral line) or 10 mL intravenous (central line) before and after infusion, or as needed for line patency Heparin 10 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Heparin 100 units/mL (#3mL or #5mL) use as a final flush for central line (QS) Flushing orders not needed  Concurrent Medications  Drug/Non-Drug Allergies  Print  Date  their representatives on my behalf to convey this prescription described herein to the and that I may not delegate signature authority.

PRESCRIBER\_

Date\_



## Patient Enrollment Form Fax completed forms to: 1-855-423-5757

RUCONEST®
Clesterase inhibitor (recombinant)

4 Patient Information

Attach copy of demographic/face she	•		DOD
Name		_	DOB
Check Preferred Phone # Work #		Home #	Cell #
Preferred Language		Caregiver Information	
mail		Caregiver Name (first, last)	
Address		Relationship to Patient	
		Caregiver Phone #	Okay to leave vm
City/State/ZIP		Caregiver Email	
5. Patient Insurance Information			
Attach copies of front and back of all	medical and p	rescription insurance cards OR o	complete below
Medical Insurance Card		<b>Prescription Drug Card</b>	
Plan Name	P	BM/Plan Name	
Plan Phone #	PI	lan Phone #	
Policy Holder Name	M	Member ID #	
Nember ID #	B	BIN #	
Group #	P	CN #	Group #
лоцр п			
6. Prescriber Information			
•	y GI Im	nmunology Primary Care Otl	ner
6. Prescriber Information		5	
6. Prescriber Information Provider Specialty: Allergy Dermatolog		NPI #	TIN #
6. Prescriber Information Provider Specialty: Allergy Dermatolog		NPI #State License #	TIN #
6. Prescriber Information Provider Specialty: Allergy Dermatolog Provider Name Medicaid Provider ID #		NPI #State License #  Office Contact Information	TIN #
6. Prescriber Information Provider Specialty: Allergy Dermatolog Provider Name Medicaid Provider ID # Site Name		NPI #  State License #  Office Contact Information  Contact Name	TIN # PTAN #
6. Prescriber Information Provider Specialty: Allergy Dermatolog Provider Name Medicaid Provider ID # Site Name		NPI #  State License #  Office Contact Information  Contact Name	TIN # PTAN #
6. Prescriber Information Provider Specialty: Allergy Dermatolog Provider Name Medicaid Provider ID # Site Name		NPI # State License # Office Contact Information Contact Name Role	TIN # PTAN #
6. Prescriber Information Provider Specialty: Allergy Dermatolog Provider Name Medicaid Provider ID # Site Name Address Sity/State/ZIP		NPI # State License # Office Contact Information  Contact Name  Role  Contact Phone	TIN # PTAN #

Before prescribing RUCONEST, please read the accompanying full Prescribing Information or go to www.ruconest.com

RUCONEST Solutions: 1-855-613-4423